# antibodies - online.com







## anti-RDH13 antibody (AA 101-200) (Alexa Fluor 555)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	RDH13
Binding Specificity:	AA 101-200
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RDH13 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human RDH13	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse,Rabbit	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	RDH13
Alternative Name:	RDH13 (RDH13 Products)

### **Target Details**

Background:	Synonyms: RDH13, RDH13_HUMAN, Retinol dehydrogenase 13 all trans and 9 cis, Retinol		
	dehydrogenase 13 all trans/9 cis, Retinol dehydrogenase 13.		
	Background: RDH13, also known as all-trans and 9-cis retinol dehydrogenase 13 or SDR7C3, is		
	a 331 amino acid mitochondrial protein belonging to the short-chain		
	dehydrogenases/reductases (SDR) family. Widely expressed, mostly in eye, pancreas, placenta		
	and lung, RDH13 localizes on the outer side of the inner mitochondrial membrane. Related to		
	microsomal retinoid oxidoreductase RDH11, RDH13 is considered to be a major enzyme		
	among the RDH family of proteins. Catalytically active, RDH13 recognizes retinoids as		
	substrates and may function in retinoic acid production. RDH13 may function to protect the		
	mitochondria against oxidative stress. Leber congenital amaurosis (LCA) type 3, an inherited		
	autosomal recessive retinal disease, has been associated with defects of RDH13. LCA		
	represents the most common genetic cause of congenital visual impairment in infants and		
	children.		
Gene ID:	112724		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		