antibodies -online.com





Datasheet for ABIN1398511

anti-PANK3 antibody (AA 271-370) (Biotin)



()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μL
Target:	PANK3
Binding Specificity:	AA 271-370
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PANK3 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PANK3
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	PANK3
Alternative Name:	PANK3 (PANK3 Products)

Target Details

Background:	Synonyms: hPanK3, MGC16863, OTTHUMP00000160961, Pank3, PANK3_HUMAN,		
· ·	Pantothenate kinase 3, pantothenic acid kinase, Pantothenic acid kinase 3.		
	Background: The pantothenate kinase (PANK) family of proteins catalyzes the first step in		
	coenzyme A (CoA) biosynthesis. Coenzyme A is an important coenzyme involved in the		
	synthesis and oxidation of fatty acids, as well as the oxidation of pyruvate in the citric acid		
	(Krebs) cycle. Pantothenate kinase 3 (PANK3) is a 370 amino acid member of the pantothenate		
	kinase family that plays a role in the physiological regulation of the intracellular CoA		
	concentration. Localized to the cytoplasm, PANK3 is regulated by feedback inhibition by CoA		
	and its thioesters. PANK3 transfers a phosphate from ATP to pantothenate (Vitamin B5),		
	resulting in formation of 4?-phosphopantothenate. Closely related to its family members,		
	PANK1, PANK2 and PANK4, PANK3 is highly expressed in liver. Pantothenate kinase associated		
	neurodegeneration (PKAN) results from mutations in the gene encoding PANK2, the only		
	mitochondria targeted human PANK.		
Gene ID:	79646		
Pathways:	Ribonucleoside Biosynthetic Process		
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		

ш	land	linc	٩
	ıanıu	11111	J

Expiry Date:

12 months