

[Go to Product page](#)

Datasheet for ABIN1398565
anti-UBE2G2 antibody (AA 100-165) (Biotin)

Overview

Quantity:	100 µL
Target:	UBE2G2
Binding Specificity:	AA 100-165
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBE2G2 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Ube2G2/UBC7
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UBE2G2
Alternative Name:	Ube2G2 (UBE2G2 Products)
Background:	Synonyms: Human ubiquitin conjugating enzyme G29, UB2G2_HUMAN, UBC 7, UBC7, UBC7

Target Details

homolog yeast, UBE2G2, Ubiquitin carrier protein G2, Ubiquitin conjugating enzyme 7, Ubiquitin conjugating enzyme E2 G2, Ubiquitin conjugating enzyme E2G 2 homologous to yeast UBC7, Ubiquitin conjugating enzyme E2G 2 UBC7 homolog yeast, Ubiquitin conjugating enzyme E2G 2, Ubiquitin conjugating enzyme G2, Ubiquitin protein ligase G2, Ubiquitin-conjugating enzyme E2 G2, Ubiquitin-protein ligase G2.

Background: UBE2G2 (Ubiquitin-conjugating enzyme E2 G2), also known as UBC7, is a 165 amino acid protein involved in ubiquitin-mediated protein degradation. Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). UBE2G2 is an E2 ubiquitin-conjugating enzyme that acts to catalyze the covalent attachment of ubiquitins to various proteins. Expressed throughout the body, UBE2G2 shares 100 % sequence identity with its mouse counterpart and is thought to be involved in endoplasmic reticulum-associated degradation (ERAD). Two isoforms of UBE2G2 exist due to alternative splicing events.

Gene ID: 7327

Application Details

Application Notes: IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Handling

Expiry Date: 12 months