

[Go to Product page](#)

Datasheet for ABIN1398688

anti-UBE2H antibody (AA 101-183) (Alexa Fluor 488)

Overview

Quantity:	100 µL
Target:	UBE2H
Binding Specificity:	AA 101-183
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBE2H antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Ube2H/UBCH2
Isotype:	IgG
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	UBE2H
Alternative Name:	Ube2H (UBE2H Products)
Background:	Synonyms: E2 20K, Ubc H2, UBC8, UBCH 2, UBCH, UBCH2, Ube2h, UBE2H_HUMAN, Ubiquitin

Target Details

carrier protein H, Ubiquitin conjugating enzyme E2 H, Ubiquitin conjugating enzyme E2H UBC8 homolog yeast, Ubiquitin conjugating enzyme E2H, Ubiquitin protein ligase H, Ubiquitin-conjugating enzyme E2 H, Ubiquitin-conjugating enzyme E2-20K, Ubiquitin-protein ligase H. Background: UBE2H, also known as UBC8, UBCH, UBCH2 or E2-20K, is a 183 amino acid protein involved in ubiquitin-mediated protein degradation. Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). One of several members of the ubiquitin-conjugating enzyme family, UBE2H functions as an E2 ubiquitin-conjugating enzyme that acts to catalyze the covalent attachment of ubiquitin residues to various proteins, including Histone H2A. UBE2H shares 100 % identity with its mouse counterpart and 98 % identity with its frog and zebrafish homologs, suggesting a conserved function between species. Multiple isoforms of UBE2H exist due to alternative splicing events.

Gene ID: 7328

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months