# antibodies - online.com







# anti-UBE2Q1 antibody (AA 321-422) (Biotin)



( )	1 /	0	rv	/ 1 /	71	Α.
	1//	$\vdash$	1 \/	16		1/1/
$\sim$	v	$\sim$	1 V	١,	_	v v

Quantity:	100 μL
Target:	UBE2Q1
Binding Specificity:	AA 321-422
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBE2Q1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human UBE2Q1
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	UBE2Q1
Alternative Name:	UBE2Q1 (UBE2Q1 Products)

## **Target Details**

Expiry Date:

rarget Details	
Background:	Synonyms: Galactosyl transferase associated protein, GTAP, NICE 5, NICE5, PRO3094, Protein
	NICE 5, Protein NICE-5, UB2Q1_HUMAN, UBE2Q, Ube2q1, Ubiquitin carrier protein Q1, Ubiquitin
	conjugating enzyme E2 Q1, Ubiquitin conjugating enzyme E2Q putative 1, Ubiquitin conjugating
	enzyme E2Q 1, Ubiquitin conjugating enzyme E2Q family member 1, Ubiquitin protein ligase Q1,
	Ubiquitin-conjugating enzyme E2 Q1, Ubiquitin-protein ligase Q1.
	Background: Ubiquitination involves at least three classes of enzymes: ubiquitin-activating
	enzymes (UBE1s), ubiquitin-conjugating enzymes (UBE2s), and ubiquitin-protein ligases
	(UBE3s). When ubiquitin is activated by a UBE1, it is transferred to the cysteine residue on a
	UBE2. UBE2 then binds a UBE3, which transfers the ubiquitin from the UBE2 cysteine to a lysine
	residue on the target protein. Ubiquitin-conjugating enzyme E2 Q1 (UBE2Q1), also known as
	ubiquitin-protein ligase Q1 or ubiquitin carrier protein Q1, is an 422 amino acid protein
	belonging to the ubiquitin-conjugating enzyme (UBE2) family. Two named isoforms of UBE2Q1
	exist as a result of alternative splicing events.
Gene ID:	55585
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

12 months