antibodies.com

Datasheet for ABIN1398778 anti-UBE2W antibody (AA 71-151) (Alexa Fluor 488)



Overview

Quantity:	100 µL	
Target:	UBE2W	
Binding Specificity:	AA 71-151	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UBE2W antibody is conjugated to Alexa Fluor 488	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human UBE2W	
Isotype:	lgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	UBE2W
Alternative Name:	UBE2W (UBE2W Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1398778 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Target Details		
Background:	Synonyms: hUBC 16, hUBC16, Probable ubiquitin conjugating enzyme E2 W, Probable ubiquitin- conjugating enzyme E2 W, UBE 2W, ube2w, UBE2W_HUMAN, Ubiquitin carrier protein W, Ubiquitin conjugating enzyme 16, Ubiquitin conjugating enzyme E2 W, Ubiquitin conjugating enzyme E2W, Ubiquitin protein ligase W, Ubiquitin-protein ligase W, FLJ11011. Background: Ubiquitin is an abundant, highly conserved protein found in all eukaryotic cells either free or covalently attached to cellular proteins. Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin- protein ligases (E3s). As an E2 class enzyme, UBE2W (Ubiquitin-conjugating enzyme E2 W), also known as Ubiquitin carrier protein W, is a 151 amino acid that catalyzes the conjugation of ubiquitin to proteins that are meant for lysosomal degradation. Functioning as a homodimer, UBE2W is widely expressed, with highest levels in testis. There are two isoforms of UBE2W that	
	are produced as a result of alternative splicing events.	
Gene ID:	55284	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1398778 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

1.1	(1:
Н	land	ling
		3

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN1398778 | 03/08/2024 | Copyright antibodies-online. All rights reserved.