

Datasheet for ABIN1398814

anti-RBX1 antibody (AA 1-70) (AbBy Fluor® 488)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	RBX1
Binding Specificity:	AA 1-70
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RBX1 antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ROC1/RNF75
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Cow
Purification:	Purified by Protein A.

Target Details

Target:	RBX1
Alternative Name:	RNF75 (RBX1 Products)

Target Details

Background: Synonyms: E3 ubiquitin-protein ligase RBX1, MGC13357, MGC1481, Protein ZYP, Rbx 1, Rbx1, RBX1_HUMAN, Regulator of Cullins 1, Ring box 1, RING box protein 1, RING finger protein 75, RING finger protein, RING-box protein 1, Ringbox protein 1, RNF 75, RNF75, ROC 1, ZYP protein. Background: E3 ubiquitin ligase component of multiple cullin-RING-based E3 ubiquitin-protein ligase complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins, including proteins involved in cell cycle progression, signal transduction, transcription and transcription-coupled nucleotide excision repair. The functional specificity of the E3 ubiquitin-protein ligase complexes depends on the variable substrate recognition components. As a component of the CSA complex promotes the ubiquitination of ERCC6 resulting in proteasomal degradation. Through the RING-type zinc finger, seems to recruit the E2 ubiquitination enzyme, like CDC34, to the complex and brings it into close proximity to the substrate. Probably also stimulates CDC34 autoubiquitination. May be required for histone H3 and histone H4 ubiquitination in response to ultraviolet and for subsequent DNA repair. Promotes the neddylation of CUL1, CUL2, CUL4 and CUL4 via its interaction with UBE2M.

Gene ID: 9978

Pathways: [Cell Division Cycle, M Phase](#), [SARS-CoV-2 Protein Interactome](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months