

Datasheet for ABIN1398815

Overview

anti-RBX1 antibody (AA 1-70) (AbBy Fluor® 555)



Go to Product page

| Quantity: | 100 μL | |
|----------------------|--------------|--|
| Target: | RBX1 | |
| Binding Specificity: | AA 1-70 | |
| Reactivity: | Human, Mouse | |
| Host: | Rabbit | |

|--|

This RBX1 antibody is conjugated to AbBy Fluor® 555

Application: Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence

(Paraffin-embedded Sections) (IF (p))

Product Details

Conjugate:

| Immunogen: | KLH conjugated synthetic peptide derived from human ROC1/RNF75 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat,Cow |
| Purification: | Purified by Protein A. |

Target Details

| Target: | RBX1 |
|-------------------|-----------------------|
| Alternative Name: | RNF75 (RBX1 Products) |

Target Details

| Background: | Synonyms: E3 ubiquitin-protein ligase RBX1, MGC13357, MGC1481, Protein ZYP, Rbx 1, Rbx1, |
|---------------------|---|
| | RBX1_HUMAN, Regulator of Cullins 1, Ring box 1, RING box protein 1, RING finger protein 75, |
| | RING finger protein, RING-box protein 1, Ringbox protein 1, RNF 75, RNF75, ROC 1, ZYP protein. |
| | Background: E3 ubiquitin ligase component of multiple cullin-RING-based E3 ubiquitin-protein |
| | ligase complexes which mediate the ubiquitination and subsequent proteasomal degradation |
| | of target proteins, including proteins involved in cell cycle progression, signal transduction, |
| | transcription and transcription-coupled nucleotide excision repair. The functional specificity of |
| | the E3 ubiquitin-protein ligase complexes depends on the variable substrate recognition |
| | components. As a component of the CSA complex promotes the ubiquitination of ERCC6 |
| | resulting in proteasomal degradation. Through the RING-type zinc finger, seems to recruit the |
| | E2 ubiquitination enzyme, like CDC34, to the complex and brings it into close proximity to the |
| | substrate. Probably also stimulates CDC34 autoubiquitination. May be required for histone H3 |
| | and histone H4 ubiquitination in response to ultraviolet and for subsequent DNA repair. |
| | Promotes the neddylation of CUL1, CUL2, CUL4 and CUL4 via its interaction with UBE2M. |
| Gene ID: | 9978 |
| Pathways: | Cell Division Cycle, M Phase, SARS-CoV-2 Protein Interactome |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |