

Datasheet for ABIN1398835
anti-FBXW7 antibody (AA 501-600) (Biotin)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FBXW7
Binding Specificity:	AA 501-600
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FBXW7 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Fbxw7/hCDC4
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	FBXW7
Alternative Name:	Fbxw7 (FBXW7 Products)

Target Details

Background:	Synonyms: AGO, CDC4, FBW6, FBW7, hAgo, FBX30, FBXW6, SEL10, hCdc4, FBXO30, SEL-10, F-box/WD repeat-containing protein 7, Archipelago homolog, F-box and WD-40 domain-containing protein 7, F-box protein FBX30, FBXW7 Background: Substrate recognition component of an SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Recognizes and binds phosphorylated sites/phosphodegrons within target proteins and thereafter bring them to the SCF complex for ubiquitination (PubMed:17434132). Identified substrates include cyclin-E (CCNE1 or CCNE2), JUN, MYC, NOTCH1 released notch intracellular domain (NICD), and probably PSEN1 (PubMed:11565034, PubMed:12354302, PubMed:11585921, PubMed:15103331, PubMed:14739463, PubMed:17558397, PubMed:17873522, PubMed:22608923). Acts as a negative regulator of JNK signaling by binding to phosphorylated JUN and promoting its ubiquitination and subsequent degradation (PubMed:14739463).
Gene ID:	55294
UniProt:	Q969H0
Pathways:	Notch Signaling , EGFR Signaling Pathway

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C for 12 months.
------------------	-------------------------------

Expiry Date:	12 months
--------------	-----------