

[Go to Product page](#)

Datasheet for ABIN1398932

anti-GLIPR2 antibody (AA 51-154) (FITC)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | GLIPR2 |
| Binding Specificity: | AA 51-154 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GLIPR2 antibody is conjugated to FITC |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human GLIPR2 |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | GLIPR2 |
| Alternative Name: | GLIPR2 (GLIPR2 Products) |
| Background: | Synonyms: GAPR 1, GAPR-1, GAPR1_HUMAN, GLI pathogenesis related 2, Glioma pathogenesis |

Target Details

related protein 2, Glioma pathogenesis-related protein 2, GliPR 2, GLIPR2, Golgi associated plant pathogenesis related protein 1, Golgi associated PR 1 protein, Golgi-associated plant pathogenesis-related protein 1, Golgi-associated PR-1 protein, OTTMUSP00000007558, RP11-421H8.5, RP23-209M8.2, C77180, 5730414A08Rik, C9orf19.

Background: Cysteine-rich secretory proteins (CRISPs) represent a family of evolutionarily conserved proteins that may play a role in the innate immune system and are transcriptionally regulated by androgens in several tissues. GAPR-1 (Golgi-associated plant pathogenesis-related protein 1), also known as GLIPR2, is a 154 amino acid lipid anchor protein belonging to the CRISP family. GAPR-1 also shares similarity with the pathogenesis-related protein (PR) superfamily, and may play an important role in the immune system. Existing as a homodimer, GAPR-1 is highly expressed in lung and peripheral leukocytes with minor expression in liver and kidney. Containing a conserved sperm-coating protein (SCP) domain, GAPR-1 binds to negatively charged lipids and may be involved in the differentiation of epithelial cells into mesenchymal cells. Increased expression of GAPR-1 in kidney may contribute to the development of fibrosis.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months