antibodies -online.com





anti-HAGHL antibody (AA 1-100) (Alexa Fluor 488)



| () | 1 / | \sim | rv | 11/ | 11 | Α |
|-----|-----|--------|------|-----|----|----|
| | 1// | ┙ | I \/ | 16 | ٦, | /\ |
| | | | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | HAGHL | |
| Binding Specificity: | AA 1-100 | |
| Reactivity: | Human, Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This HAGHL antibody is conjugated to Alexa Fluor 488 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human HAGHL | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse | |
| Predicted Reactivity: | Rat,Dog,Cow,Sheep,Pig,Horse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | HAGHL | |
|-------------------|------------------------|--|
| Alternative Name: | HAGHL (HAGHL Products) | |

Target Details

Storage Comment:

Expiry Date:

| rarget Details | |
|---------------------|--|
| Background: | Synonyms: HAGHL, HAGHL_HUMAN, Hydroxyacylglutathione hydrolase-like, |
| | Hydroxyacylglutathione hydrolase-like protein. |
| | Background: The hydroxyacylglutathione hydrolase-like protein (HAGHL) is a 290 amino acid |
| | protein that belongs to the glyoxalase II family. HAGHL binds two zinc ions per subunit and acts |
| | as a hydrolase on ester bonds. The gene encoding HAGHL maps to human chromosome 16, |
| | which encodes over 900 genes in approximately 90 million base pairs, makes up nearly 3 % of |
| | human cellular DNA and is associated with a variety of genetic disorders. The rare disorder |
| | Rubinstein-Taybi syndrome is associated with chromosome 16, as is Crohn's disease, a |
| | gastrointestinal inflammatory condition that may involve the NOD2 gene. An association with |
| | systemic lupus erythematosis and a number of other autoimmune disorders with the |
| | pericentromeric region of chromosome 16 has led to the identification of SLC5A11 as a |
| | potential autoimmune modifier. |
| Gene ID: | 84264 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| 01 0 1 | |

12 months

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.