

[Go to Product page](#)

Datasheet for ABIN1399323

anti-FNDC3B antibody (AA 921-1020) (Alexa Fluor 350)

Overview

Quantity:	100 µL
Target:	FNDC3B
Binding Specificity:	AA 921-1020
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FNDC3B antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FNDC3B
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Sheep, Pig, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FNDC3B
Alternative Name:	FNDC3B (FNDC3B Products)
Background:	Synonyms: Factor for adipocyte differentiation 104, FAD104, fibronectin type III domain

Target Details

containing 3B, FLJ23399, HCV NS5A binding protein 37, NS5ABP37, YVTM2421, MGC10002, PRO4979, DKFZp686D14170, DKFZp762K137, FND3B_HUMAN.

Background: Adipogenesis, the process of transforming pre-adipocytes into mature fat cells, is of particular interest due to the role adipocytes play in obesity and type II diabetes. Adipocytes have been shown to affect a variety of functions, including hemostasis, angiogenesis and energy balance, by secreting hormones and bioactive peptides. The FNDC3B protein, also designated FAD104 (factor for adipocyte differentiation 104) or HCV NS5A-binding protein 37, is expressed during early adipogenesis. Belonging to the FNDC3 family of proteins, FNDC3B is a 1,204 amino acid protein that contains nine fibronectin type-III domains. FNDC3B-deficient mice die within one day of birth, suggesting that FNDC3B is crucial for postpartum survival. Mouse embryonic fibroblasts (MEFs) with loss of FNDC3B function displayed a reduction in stress fiber formation, indicating a role for FNDC3B in cell proliferation, adhesion, spreading and migration.

Gene ID: 64778

Pathways: [Positive Regulation of fat Cell Differentiation](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months