

Datasheet for ABIN1399494

anti-ELL antibody (AA 531-620) (AbBy Fluor® 647)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ELL
Binding Specificity:	AA 531-620
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ELL antibody is conjugated to AbBy Fluor® 647
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ELL
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Horse
Purification:	Purified by Protein A.

Target Details

Target:	ELL
Alternative Name:	ELL (ELL Products)
Background:	Synonyms: ELL_HUMAN, Eleven nineteen lysine rich leukemia gene, Eleven-nineteen lysine-rich

Target Details

leukemia protein, ELL1, elongation factor RNA polymerase II, Men, RNA polymerase II elongation factor, RNA polymerase II elongation factor ELL.

Background: Eukaryotic RNA polymerase II mediates the synthesis of mature and functional messenger RNA. This is a multistep process, called the transcription cycle, that includes five stages: preinitiation, promoter, clearance, elongation and termination. Elongation is thought to be a critical stage for the regulation of gene expression. ELL (11-19 lysine-rich leukemia protein), also designated MEN, functions as an RNA polymerase II elongation factor that increases the rate of transcription by suppressing transient pausing by RNA polymerase II. It is also thought to regulate cellular proliferation. ELL is abundantly expressed in peripheral blood leukocytes, skeletal muscle, placenta and testis, with lower expression in spleen, thymus, heart, brain, lung, kidney, liver and ovary. The gene encoding human ELL, which maps to chromosome 19p13.1, is one of several genes that undergo translocation with the MLL gene on chromosome 11q23 in acute myeloid leukemia. MLL (myeloid/lymphoid leukemia, also designated ALL-1 and HRX) regulates embryonal and hematopoietic development.

Gene ID: 8178

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months