

[Go to Product page](#)

Datasheet for ABIN1399567  
**anti-PPAP2C antibody (AA 151-250) (Biotin)**

## Overview

Quantity:	100 µL
Target:	PPAP2C
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPAP2C antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human VANGL1/LPP2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	PPAP2C
Alternative Name:	LPP2 ( <a href="#">PPAP2C Products</a> )
Background:	Synonyms: Loop tail protein 2 homolog, Loop-tail protein 2 homolog, LPP2, MGC5338, STB2,

## Target Details

STBM2, Strabismus 2, Van Gogh like protein 1, Van Gogh-like protein 1, vang like 1 van gogh, Drosophila, Vang like protein 1, Vang-like protein 1, VANG1\_HUMAN, VANGL 1, Vangl1.

Background: The Vang family of proteins are integral membrane proteins that are homologues of the Drosophila tissue polarity gene strabismus. The gene encoding for Van Gogh-like protein 1 (Vangl1), also designated Strabismus 2 (STB2), localizes to human chromosome 1p11-p13.1. Van Gogh-like protein 2 (Vangl2), also designated Strabismus 1 (STB1), localizes to chromosome 1q22-q23. Vangl1 is expressed primarily in testis and ovary, but is also expressed in gastric and pancreatic cancer. Vangl proteins play a key developmental role in establishing planar cell polarity (PCP) and in regulating convergent extension (CE) movements during embryogenesis. Vangl1 and Vangl2 are both downregulated in several cancer cell lines and primary tumors.

Gene ID: 8612

## Application Details

Application Notes: WB 1:300-5000  
IHC-P 1:200-400  
IHC-F 1:100-500

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months