antibodies - online.com







anti-MLLT11 antibody (AA 41-90) (FITC)



| () | 1 / | \sim | KI / | 110 | Νę |
|-----|-----|----------|------|-----|-------|
| | 1// | \vdash | I \/ | 1 ← | ٠// ٢ |
| | | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | MLLT11 | |
| Binding Specificity: | AA 41-90 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This MLLT11 antibody is conjugated to FITC | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human MLLT11 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | MLLT11 |
|-------------------|--------------------------|
| Alternative Name: | MLLT11 (MLLT11 Products) |

Target Details

| Bacl | // | ro | ıın | ٨. |
|------|----|----|-----|----|
| Daci | ĸα | ΙU | uu | u. |

Synonyms: AF1Q, ALL1 fused gene from chromosome 1q, MLLT 11, MLLT-11,

Myeloid/lymphoid or mixed lineage leukemia trithorax homolog, Drosophila translocated to 11, Protein AF1q, RP11 316M1.10.

Background: The gene encoding the Mixed-Lineage Leukemia (MLL) proteins is located on chromosome 11q23. Chromosomal translocations involving band 11q23 result in rogue activator proteins that are associated with approximately 10 % of patients with acute lymphoblastic leukemia (ALL) and 5 % of patients with acute myeloid leukemia (AML). Most patients affected are less than 1 year of age. MLLT11, also known as mixed-lineage leukemia translocated to 11 or AF1q, is a 90 amino acid MLL fusion partner. Based on the expression patterns of MLLT11, it is thought that MLLT11 plays a role in leukemogenesis and, specifically, the progression of acute monocytic leukemia (AML). Also, expressed in embryonic brain cortex, MLLT11 is upregulated during neuronal differentiation and is thought to play a role in the development of the central nervous system. Finally, MLLT11 has been shown to be differentially expressed in highly metastatic cells, in comparison with non-metastatic parent cells. Such findings suggest a role of MLLT11 in tumorigenesis.

Gene ID:

10962

Application Details

| An | plication | n Notes |
|------------|-----------|---------|
| $\neg \nu$ | piicatioi | TINOLCS |

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |