antibodies -online.com





anti-FSIP1 antibody (AA 101-200) (Alexa Fluor 488)



Overview

Alternative Name:

| Quantity: | 100 μL |
|-----------------------|--|
| Target: | FSIP1 |
| Binding Specificity: | AA 101-200 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FSIP1 antibody is conjugated to Alexa Fluor 488 |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), |
| | Immunofluorescence (Cultured Cells) (IF (cc)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human FSIP1 |
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human,Mouse,Dog,Cow,Sheep |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | FSIP1 |

FSIP1 (FSIP1 Products)

Target Details

| Background: | Synonyms: Fibrous sheath interacting protein 1, FLJ35989, HSD10, FSIP1_HUMAN. |
|---------------------|---|
| | Background: FSIP1 is a 581 amino acid protein that is expressed in airway epithelium. A |
| | member of the FSIP1 family, FSIP1 is encoded by a gene that maps to human chromosome |
| | 15q14 and mouse chromosome 2 E5. Encoding more than 700 genes, chromosome 15 is |
| | made up of approximately 106 million base pairs and is about 3 % of the human genome. |
| | Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes |
| | in the 15q11-q13 region. In the case of Angelman syndrome, this loss is due to inactivity of the |
| | maternal 15q11-q13 encoded UBE3A gene in the brain by either chromosomal deletion or |
| | mutation. In cases of Prader-Willi syndrome, there is a partial or complete deletion of this region |
| | from the paternal copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated |
| | with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is |
| | associated with chromosome 15 through the FBN1 gene. |
| Gene ID: | 161835 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |