antibodies -online.com





anti-HLA-B27 antibody (AA 81-180) (Biotin)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	HLA-B27
Binding Specificity:	AA 81-180
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HLA-B27 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HLA B27	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

Target Details

Target:	HLA-B27
Alternative Name:	HLA B27 (HLA-B27 Products)
Background: Synonyms: HLA B27 HLA-B27, HLAB, Leukocyte antigen class I B, Lymphocyte antigen,	

histocompatibility complex class I B, SPDA1, 1B27_HUMAN.

Background: Major histocompatibility complex (MHC) molecules form an integral part of the immune response system. They are cell-surface receptors that bind peptides and present them to T lymphocytes. Human leukocyte antigens (HLAs) are polymorphic members of the MHC family that are specifically involved in the presentation of antigens to the T cell receptor. There are two classes of HLA antigens: class I (HLA-A, HLA-B and HLA-C) and class II (HLA-D). Class I molecules are expressed in nearly all cells and play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum. The differential structural properties of MHC class I and class II molecules account for their respective roles in activating different populations of T lymphocytes. HLA-B encodes a membrane anchored heavy chain which hetero-dimerizes with a light chain (-2-Microglobulin) to form MHC-I. Polymorphisms yield hundreds of HLA-B alleles. The HLA-B27 allele appears with increased frequency in uveitis patients.

Application Details

WB 1:300-5000

Application Notes:

Application Notes.	WB 1.500-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months