antibodies -online.com





anti-TAS2R38 antibody (AA 151-250) (Alexa Fluor 555)



Overview

| Quantity: | 100 μL |
|----------------------|---|
| Target: | TAS2R38 |
| Binding Specificity: | AA 151-250 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TAS2R38 antibody is conjugated to Alexa Fluor 555 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human T2R38/TAS2R38 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Chicken |
| Purification: | Purified by Protein A. |

Target Details

| Target: | TAS2R38 |
|-------------------|--------------------------|
| Alternative Name: | T2R38 (TAS2R38 Products) |

Target Details

| | 0 |
|---------------------|---|
| Background: | Synonyms: PTC, PTC bitter taste receptor, T2R38, T2R61, taste receptor, type 2, member 38, |
| | T2R38_HUMAN. |
| | Background: The sense of taste is essential for the survival of organisms. For example, the |
| | ability to identify sweet-tasting foods enables animals to seek out food with high nutritive value, |
| | whereas the ability to identify bitter substances enables them to avoid the ingestion of |
| | potentially harmful substances. A family of integral membrane proteins are involved in taste |
| | perception and include T1R, which is involved in sweet taste perception and T2R, which is |
| | involved in bitter taste perception. Both types of taste receptors couple to various G proteins to |
| | initiate signal transduction cascades. Specifically, T2R38 is expressed in subsets of taste |
| | receptor cells of the tongue and exclusively in gustducin-positive cells. Variations in T2R38 are |
| | associated with the ability to taste the bitter chemical phenylthiocarbamide (PTC), also called |
| | thiourea tasting. |
| Gene ID: | 5726 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| , ppcation retor | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |