# antibodies -online.com





#### Datasheet for ABIN1400081

# anti-BASP1 antibody (AA 3-100) (AbBy Fluor® 555)



| ( ) | 11/0   | K\ /   | iew         |
|-----|--------|--------|-------------|
|     | $\cup$ | 'I V/I | $I \cap VV$ |
|     |        |        |             |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | BASP1   |  |
| Binding Specificity: | AA 3-100  |  |
| Reactivity:          | Human, Mouse, Rat   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This BASP1 antibody is conjugated to AbBy Fluor® 555  |  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human BASP1/Nap22 |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Cross-Reactivity:     | Human, Mouse, Rat   |  |
| Predicted Reactivity: | Dog,Cow,Pig,Horse,Chicken,Rabbit                                |  |
| Purification:         | Purified by Protein A.  |  |

## **Target Details**

| Target:           | BASP1                  |
|-------------------|------------------------|
| Alternative Name: | BASP1 (BASP1 Products) |

## **Target Details**

| _                   |   |  |
|---------------------|---|--|
| Background:         | Synonyms: CAP23, NAP22, CAP-23, NAP-22, Brain acid soluble protein 1, 22 kDa neuronal           |  |
|                     | tissue-enriched acidic protein, Neuronal axonal membrane protein NAP-22, BASP1                  |  |
|                     | Background: Neuronal axonal membrane protein Nap-22, also designated neuronal tissue-           |  |
|                     | enriched acidic protein or brain acid soluble protein (BASP1), is a Ca2+-dependent calmodulin-  |  |
|                     | binding protein that is important for neuronal sprouting and plasticity. Nap-22 is abundant in  |  |
|                     | brain nerve terminals and is also present in significant amounts in kidney, testis and lymphoid |  |
|                     | tissue. Nap-22 undergoes N-terminal myristoylation for membrane localization. It has been       |  |
|                     | characterized as a major protein of neuronal rafts, which are known to preferentially bind      |  |
|                     | membranes containing cholesterol. Nap-22 is a crucial protein active in neurite outgrowth and   |  |
|                     | synaptic plasticity.  |  |
| Gene ID:            | 10409   |  |
| UniProt:            | P80723  |  |
| Application Details |   |  |
| Application Notes:  | IF(IHC-P) 1:50-200  |  |
|                     | IF(IHC-F) 1:50-200  |  |
|                     | IF(ICC) 1:50-200  |  |
| Restrictions:       | For Research Use only   |  |
| Handling            |   |  |
| Format:             | Liquid  |  |
| Concentration:      | 1 μg/μL   |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and     |  |
|                     | 50 % Glycerol.  |  |
| Preservative:       | ProClin   |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be             |  |
|                     | handled by trained staff only.  |  |
| Storage:            | -20 °C  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.               |  |
| Expiry Date:        | 12 months   |  |
|                     |   |  |