## antibodies .- online.com





## anti-SARM1 antibody (AA 340-380) (Biotin)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	SARM1
Binding Specificity:	AA 340-380
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SARM1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SARM1	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

## Target Details

Target:	SARM1	
Alternative Name:	SARM1 (SARM1 Products)	
Background:	Synonyms: SARM, SAMD2, MyD88-5, Sterile alpha and TIR motif-containing protein 1, Sterile	
	alpha and Armadillo repeat protein, Sterile alpha motif domain-containing protein 2, SAM	

	domain-containing protein 2, Tir-1 homolog, SARM1, KIAA0524	
	Background: Negative regulator of MYD88- and TRIF-dependent toll-like receptor signaling	
	pathway which plays a pivotal role in activating axonal degeneration following injury. Promotes	
	Wallerian degeneration an injury-induced axonal death pathway which involves degeneration of	
	an axon distal to the injury site. Can activate neuronal death in response to stress. Regulates	
	dendritic arborization through the MAPK4-JNK pathway. Involved in innate immune response.	
	Inhibits both TICAM1/TRIF- and MYD88-dependent activation of JUN/AP-1, TRIF-dependent	
	activation of NF-kappa-B and IRF3, and the phosphorylation of MAPK14/p38.	
Gene ID:	23098	
UniProt:	Q6SZW1	
Pathways:	TLR Signaling, Activation of Innate immune Response, Toll-Like Receptors Cascades	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	