antibodies -online.com





anti-RCBTB1 antibody (Alexa Fluor 555)



Go to Product page

| () | 11/0 | K\ / | iew | 1 |
|-----|--------|------|---------------------|---|
| | \cup | ועוי | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μL |
|--------------|---|
| Target: | RCBTB1 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RCBTB1 antibody is conjugated to Alexa Fluor 555 |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human RCBTB1/CLLD7 |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | RCBTB1 | |
|-------------------|---|--|
| Alternative Name: | RCBTB1 (RCBTB1 Products) | |
| Background: | Synonyms: Chronic lymphocytic leukemia deletion region gene 7 protein, CLL deletion region gene 7 protein, CLLD7, CLLL7 protein, Guanine nucleotide exchange factor, broad-complex, | |
| | tramtrack and bric a brac, Hypothetical protein FLJ39335, RCBT1_HUMAN, Rcbtb1, RCC1 and BTB domain-containing protein 1, Regulator of chromosome condensation and BTB domain- | |
| | bib domain containing protein i, regulator of chloriosome condensation and bib domain- | |

containing protein 1.

Background: The BTB domain, also known as the POZ (poxvirus and zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C2H2-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. RCBTB1 (regulator of chromosome condensation (RCC1) and BTB (POZ) domain containing protein 1), also known as GLP, CLLD7, CLLL7 or E4.5, is a 531 amino acid protein that localizes to the nucleus and contains two BTB (POZ) domains and six RCC1 repeats. Expressed ubiquitously, RCBTB1 is thought to be involved in cell cycle regulation, specifically via chromatin remodeling. The gene encoding RCBTB1 maps to a region on human chromosome 13 that is frequently deleted in B-cell chronic lymphocytic leukemia, suggesting a possible role for RCBTB1 in tumor suppression.

Gene ID:

55213

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|-----------------------|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |

Concentration: $1 \mu g/\mu L$ Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. -20 °C Storage: Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. 12 months **Expiry Date:**