



[Go to Product page](#)

Datasheet for ABIN1400390  
**anti-IQCG antibody (FITC)**

### Overview

Quantity:	100 µL
Target:	IQCG
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IQCG antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IQCG
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### Target Details

Target:	IQCG
Alternative Name:	IQCG ( <a href="#">IQCG Products</a> )
Background:	Synonyms: IQ domain containing protein G, IQ mot containing G, DKFZp434B227, FLJ11667, FLJ23571, FLJ37775, IQCG_HUMAN. Background: IQCG is a 443 amino acid protein containing one IQ domain. Widely distributed in nature, the IQ domain forms an amphiphilic seven-turn -helix capable of binding calmodulin in a

## Target Details

---

Ca<sup>2+</sup>-independent manner. The level of intracellular calcium is tightly regulated in all eukaryotic cells. A modest increase in this level can result in a myriad of physiological responses, most of which are mediated by calmodulin (CaM), the universal calcium sensor. In acute T-lymphoid/myeloid leukemia, IQCG forms a complex with Nup98, an O-linked glycoprotein and a component of the nuclear pore complex. NUP98-IQCG complex bind co-activators and/or co-repressors, which suggest a role in transcriptional regulation. Nup98-IQCG complex inhibits 32Dcl3 cell apoptosis induced by Arabinofuranosylcytosine (Ara-C) and partially blocks granulocyte differentiation induced by G-CSF. IQCG exists as two isoforms due to alternatively splicing events.

---

Gene ID: 84223

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months