# antibodies .- online.com





# anti-PDZD2 antibody (AbBy Fluor® 350)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	PDZD2	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PDZD2 antibody is conjugated to AbBy Fluor® 350	
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PDZK3	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	PDZD2	
Alternative Name:	PDZK3 (PDZD2 Products)	
Background:	Synonyms: Activated in prostate cancer protein, AIPC, KIAA0300, Likely ortholog of rat	
	plakophilin related armadillo repeat protein interacting PDZ protein antibody PAPIN, PDZ	
	domain containing 2, PDZ domain containing 3, PDZ domain containing protein 2, PDZ domain	
	containing protein 3, PDZD2, PDZK 2, PDZK2, PDZK-2, PDZK 3, PDZK3, PDZK-3, PIN1,	

#### **Target Details**

Plakophilin related armadillo repeat protein interacting PDZ protein like, Rat plakophilin related armadillo repeat protein interacting PDZ protein like, PDZD2\_HUMAN.

Background: Proteins containing PDZ domains have been shown frequently to bind the C-termini of transmembrane receptors or ion channels. They have also been shown to bind to other PDZ domain proteins and could possibly be involved in intracellular signalling. PDZK3 contains six PDZ domains and shares sequence similarity with pro-interleukin-16 (pro-IL-16). Like pro-IL-16, the encoded protein localizes to the endoplasmic reticulum and is thought to be cleaved by a caspase to produce a secreted peptide containing two PDZ domains. In addition, the PDZK3 gene is upregulated in primary prostate tumors and may be involved in the early stages of prostate tumorigenesis.

Gene ID:

23037

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	

#### Handling

r landing	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months