# antibodies .- online.com







# anti-DPCR1 antibody (Alexa Fluor 555)



( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	DPCR1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPCR1 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DPCR1	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	DPCR1	
Alternative Name:	DPCR1 (DPCR1 Products)	
Background:	Synonyms: Dfuse panbronchiolitis critical region 1, Dfuse panbronchiolitis critical region, Dfuse panbronchiolitis critical region protein 1, DKFZp6660235, DPCR protein, MGC126710,	
	MGC126712, OTTHUMP00000062447, PBLT, bCX105N19.6, DPCR1_HUMAN.	
	Background: Making up nearly 6 % of the human genome, chromosome 6 contains around	

#### **Target Details**

1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through the HFE gene which, when mutated, predisposes an individual to developing this porphyria. Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes encoding the major histocompatiblity complex proteins, which are key molecular components of the immune system and determine predisposition to rheumatic diseases, are also located on chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has been identified on the q arm of chromosome 6.

Gene ID:

135656

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months