antibodies

## Datasheet for ABIN1400586 anti-DPCR1 antibody (Alexa Fluor 647)



~	
Over	
	VIC VV

Overview	
Quantity:	100 μL
Target:	DPCR1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPCR1 antibody is conjugated to Alexa Fluor 647
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human DPCR1
lsotype:	lgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.
Target Details	
Target:	DPCR1
Alternative Name:	DPCR1 (DPCR1 Products)
Background:	Synonyms: Dfuse panbronchiolitis critical region 1, Dfuse panbronchiolitis critical region, Dfuse
	panbronchiolitis critical region protein 1, DKFZp6660235, DPCR protein, MGC126710,
	MGC126712, OTTHUMP00000062447, PBLT, bCX105N19.6, DPCR1_HUMAN.
	Background: Making up nearly 6 % of the human genome, chromosome 6 contains around

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1400586 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	1,200 genes within 170 million base pairs of sequence. Deletion of a portion of the q arm of
	chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a
	cancer susceptibility locus. Porphyria cutanea tarda is associated with chromosome 6 through
	the HFE gene which, when mutated, predisposes an individual to developing this porphyria.
	Notably, the PARK2 gene, which is associated with Parkinson's disease, and the genes
	encoding the major histocompatiblity complex proteins, which are key molecular components
	of the immune system and determine predisposition to rheumatic diseases, are also located on
	chromosome 6. Stickler syndrome, 21-hydroxylase deficiency and maple syrup urine disease
	are also associated with genes on chromosome 6. A bipolar disorder susceptibility locus has
	been identified on the q arm of chromosome 6.
Gene ID:	135656
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months