antibodies .- online.com





anti-CINP antibody (Alexa Fluor 555)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	CINP
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CINP antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human CINP
	KLH conjugated synthetic peptide derived from human CINP
lmmunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Human, Mouse, Rat
Immunogen: Isotype: Cross-Reactivity: Purification:	IgG Human, Mouse, Rat
Immunogen: Isotype: Cross-Reactivity: Purification: Target Details	IgG Human, Mouse, Rat Purified by Protein A.

kinase 2 interacting protein, Cyclin-dependent kinase 2-interacting protein, MGC849.

Background: Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin

dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key

Target Details

substrates involved in each phase of cell cycle progression. Specifically, Cdk2 interacts with Cyclins A, B1, B3, D, or E to control cell cycle progression. The Cyclin-dependent kinase 2interacting protein (CINP) interacts with components of the replication complex and Cdk2 and Cdc7, thereby providing a functional and physical link between Cdk2 and Cdc7 during firing of the origins of replication. However, CINP is phopshorylated by Cdc7, but not by Cdk2. CINP also interacts with ATR-interacting protein and regulates ATR-dependent signaling, resistance to replication stress and G2 checkpoint integrity.

Gene ID:

51550

1 μg/μL

Application Details

Concentration:

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin

Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	