.-online.com antibodies

Datasheet for ABIN1400707 anti-CINP antibody (Biotin)



Overview

Overview	
Quantity:	100 µL
Target:	CINP
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CINP antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human CINP
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	CINP
Alternative Name:	CINP (CINP Products)
Background:	Synonyms: CINP protein, CDK2-interacting protein, CINP, CINP_HUMAN, Cyclin dependent
	kinase 2 interacting protein, Cyclin-dependent kinase 2-interacting protein, MGC849.

Background: Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin

dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1400707 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

Target Details

substrates involved in each phase of cell cycle progression. Specifically, Cdk2 interacts with
Cyclins A, B1, B3, D, or E to control cell cycle progression. The Cyclin-dependent kinase 2-
interacting protein (CINP) interacts with components of the replication complex and Cdk2 and
Cdc7, thereby providing a functional and physical link between Cdk2 and Cdc7 during firing of
the origins of replication. However, CINP is phopshorylated by Cdc7, but not by Cdk2. CINP also
interacts with ATR-interacting protein and regulates ATR-dependent signaling, resistance to
replication stress and G2 checkpoint integrity.

Gene ID: 51550

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months