# antibodies - online.com







## anti-FBXW12 antibody (Biotin)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:100 μLTarget:FBXW12Reactivity:HumanHost:Rabbit	
Reactivity: Human	
<u> </u>	2
Host: Rabbit	
Clonality: Polyclor	nal
Conjugate: This FBX	XW12 antibody is conjugated to Biotin
Application: Western	Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human FBXW12/FBXO35	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	FBXW12	
Alternative Name:	FBXW12 (FBXW12 Products)	
Background:	Synonyms: F box and WD 40 domain containing protein 12; F box and WD 40 domain protein 12; F box and WD repeat domain containing 12; F box and WD40 repeat containing protein 12; F box and WD40 repeat containing 12; F box and WD40 repeat containing Protein 12; F box and WD40 repeat containin	
	box only protein 35; F box/WD repeat containing protein 12; F-box and WD-40 domain-containing protein 12; F-box only protein 35; F-box/WD repeat-containing protein 12; Fbw12;	

FBW12\_HUMAN; FBX12; FBXO12; FBXO35; FBXW 12; FBXW12.

Background: Belonging to the F-box family of proteins, FBXO35 (F-box only protein 35), also designated F-box and WD-40 domain-containing protein 12 (FBXW12 or FBXO12), is a 464 amino acid protein that contains one C-terminal F-box domain. F-box proteins are critical components of the SCF (Skp1-CUL-1-F-box protein)-type E3 ubiquitin ligase complex and are involved in substrate recognition and recruitment for ubiquitination. They are members of a larger family of proteins that are involved in the regulation of a wide variety of cellular mechanisms, including the cell cycle, the immune response, signaling cascades and developmental processes. They function by targeting proteins, such as cyclins, cyclindependent kinase inhibitors, 1°B-å and -catenin, for degradation by the proteasome after ubiquitination. Via its F-box domain, FBXO35 can directly interact with Skp1 p19 and CUL-1. FBXO35 is ubiquitously expressed at low levels in most human tissues.

Gene ID:

285231

#### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions: For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months