antibodies -online.com





anti-ATAD3A antibody (Biotin)



\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μL
Target:	ATAD3A
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATAD3A antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ATAD3A
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	ATAD3A
Alternative Name:	ATAD3A (ATAD3A Products)
Background:	Synonyms: AAA-ATPase TOB3, ATAD3B, ATAD3C, Atad3a, ATPase family AAA domain
	containing protein 3A, ATPase family AAA domain-containing protein 3, ATPase family, AAA
	domain containing 3A, FLJ10709, FLJ35514, AAA ATPase TOB3, AAA-TOB3, ATPase family
	AAA domain containing protein 3B, ATPase family, AAA domain containing 3B, ATPase family,

AAA domain containing, member 3B, TOB3, ATD3A_HUMAN, ATAD3A/B/C.

Background: The AAA ATPase family of molecular chaperones are characterized by a highly conserved AAA motif. Composed of 200-250 residues, the AAA domain contains Walker homology sequences and imparts ATPase activity. Members of the AAA ATPase family act as DNA helicases as well as transcription factors and are thought to be involved in several cellular functions such as cell-cycle regulation, protein proteolysis, organelle biogenesis and vesicle-mediated protein transport. Mitochondrial membrane proteins ATAD3A and ATAD3B contribute to the stabilization of nucleoids which are large mitochondrial DNA (mtDNA)-protein complexes. ATAD3A/B may participate in the transformation pathway and the chemosensitivity of oligodendrogliomas. The genes encoding ATAD3A/B/C maps to human chromosome 1, which houses over 3,000 genes and is the largest human chromosome spanning about 260 million base pairs and making up 8 % of the human genome.

Gene ID:

55210

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months