# antibodies -online.com





# anti-COPS4 antibody (Biotin)



Go to	Prod	luct	page
		uot	Puge

$\sim$					
	1/6	⊃r	\/I	$\triangle$	٨/

100 μL COPS4
COPS4
Human, Mouse, Rat
Rabbit
Polyclonal
This COPS4 antibody is conjugated to Biotin
Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human COPS4
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	COPS4
Alternative Name:	COPS4 (COPS4 Products)
Background:	Synonyms: CH4, COP9 constitutive photomorphogenic homolog subunit 4, COP9 complex homolog subunit 4, COP9 complex S4, COP9 constitutive photomorphogenic homolog subunit
	4, COP9 signalosome complex subunit 4, COP9 signalosome subunit 4, COPS4, CSN complex
	subunit 4, CSN4, CSN4 SCHPO, CSN4_HUMAN, DCH4, JAB1 containing signalosome subunit 4,

JAB1-containing signalosome subunit 4, MGC94896, SGN4, Signalosome subunit 4. Background: Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (UbI) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the UbI ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the UbI system, respectively.

Gene ID:

51138

Pathways:

Cell Division Cycle

### **Application Details**

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

Restrictions:

For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months