antibodies -online.com





anti-SH3RF1 antibody (Biotin)



Go to Product page

\sim					
	1/6	⊃r	\/I	\triangle	٨/

Quantity:	100 μL
Target:	SH3RF1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH3RF1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human SH3MD2/RNF142
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	SH3RF1
Alternative Name:	RNF142 (SH3RF1 Products)
Background:	Synonyms: Plenty of SH3 domains, Plenty of SH3s, POSH, Ring finger protein 142, RNF 142, RNF142, SH3 domain containing ring finger 1, SH3 domain containing RING finger protein 1, SH3MD2, SH3 multiple domains 2, SH3RF1.
	Background: SH3MD2 acts as a scaffold protein, contributes to Rac-induced signal

transduction such as JNKs (MAPK8 and MAPK9) activation and induces apoptosis. Within a signaling complex, it probably recruits protein kinases such as MAP3K10 or MAP3K11 which are in turn activated leading to the sequential activation of MAP2K4, MAP2K7 and JNKs (MAPK8 and MAPK9). SH3MD2 may be involved in targeting of HIV-1 GAG and GAG-POL polyproteins to the plasma membrane. This gene encodes a protein containing an N-terminus RING-finger, four SH3 domains, and a region implicated in binding of the Rho GTPase Rac. Via the RING-finger, the encoded protein has been shown to function as an ubiquitin-protein ligase involved in protein sorting at the trans-Golgi network. The encoded protein may also act as a scaffold for the c-Jun N-terminal kinase signaling pathway, facilitating the formation of a functional signaling module. There are two named isoforms.

Gene ID:

57630

Application Details

Application Notes: WB 1:300-5000 IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months