antibodies -online.com





anti-TRIM4 antibody (Biotin)



| \sim | |
|---------|-----------|
| ()\/户 | rview |
| \circ | V I C V V |

| Quantity: | 100 μL |
|--------------|--|
| Target: | TRIM4 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TRIM4 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |
| | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human RNF89 |
|-------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Purified by Protein A. |

Target Details

| Target: | TRIM4 |
|-------------------|--|
| Alternative Name: | RNF87 (TRIM4 Products) |
| Background: | Synonyms: Ring finger protein 87, RNF 87, TRIM4, Tripartite mot containing 4, Tripartite mot protein 4, Tripartite mot protein TRIM4, Background: The protein encoded by this gene is a member of the tripartite motif (TRIM) family. |
| | The TRIM motif includes three zinc binding domains, a RING, a B box type 1 and a B box type 2, |

Target Details

| rangerbetane | |
|---------------------|--|
| | and a coiled coil region. The protein localizes to cytoplasmic bodies. Its function has not been identified. Alternative splicing of this gene generates three transcript variants, named alpha, beta and gamma. |
| Gene ID: | 89122 |
| Application Details | |
| Application Notes: | WB 1:300-5000 IHC-P 1:200-400 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Expiry Date: | 12 months |