

Datasheet for ABIN1401257

**anti-RNF8 antibody (AbBy Fluor® 555)**[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	RNF8
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF8 antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RNF8
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## Target Details

Target:	RNF8
Alternative Name:	RNF8 ( <a href="#">RNF8 Products</a> )
Background:	<p>Synonyms: C3HC4 type zinc finger protein, E3 ubiquitin protein ligase RNF8, E3 ubiquitin-protein ligase RNF8, Ring finger protein C3HC4 type 8, RING finger protein 8, RNF 8, RNF8, RNF8_HUMAN, UBC13/UEV interacting ring finger protein.</p> <p>Background: The RING finger motif is a specialized DNA-binding zinc finger domain found in</p>

## Target Details

many transcriptional regulatory proteins. The ring finger protein (RNF) family includes any protein containing the signature RING finger motif. RNF8 is a ubiquitously expressed nuclear RING finger protein that acts as an E3 ubiquitin-protein ligase. It is required for the ubiquitination of some nuclear proteins and promotes their subsequent degradation. The heterodimeric ubiquitin-conjugating enzyme UBC13 interacts with RNF8, and they co-localize in the nucleus. RNF8 may regulate mediation of UBC13 polyubiquitylation by elongating the ubiquitin chains. RNF8 also binds to Retinoid X receptor alpha (RXR $\alpha$ ), a member of the steroid hormone receptor superfamily. It increases RXR $\alpha$ -mediated transactivation of the RXR $\alpha$ -responsive element (RXRE) promoter in a dose-dependent manner, suggesting that RNF8 is a regulator of RXR $\alpha$ -mediated transcriptional activity.

Gene ID:	9025
Pathways:	<a href="#">Production of Molecular Mediator of Immune Response</a>

## Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 $\mu$ g/ $\mu$ L
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months