# antibodies - online.com





# anti-RNF8 antibody (Biotin)



$\sim$			
	N/P	r\/	i⊢₩

0.0		
Quantity:	100 μL	
Target:	RNF8	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RNF8 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

## **Product Details**

lmmunogen:	KLH conjugated synthetic peptide derived from human RNF8	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	RNF8	
Alternative Name:	RNF8 (RNF8 Products)	
Background:	Synonyms: C3HC4 type zinc finger protein, E3 ubiquitin protein ligase RNF8, E3 ubiquitin-protein	
	ligase RNF8, Ring finger protein C3HC4 type 8, RING finger protein 8, RNF 8, RNF8,	
	RNF8_HUMAN, UBC13/UEV interacting ring finger protein.	
	Background: The RING finger motif is a specialized DNA-binding zinc finger domain found in	

many transcriptional regulatory proteins. The ring finger protein (RNF) family includes any protein containing the signature RING finger motif. RNF8 is a ubiquitously expressed nuclear RING finger protein that acts as an E3 ubiquitin-protein ligase. It is required for the ubiquitination of some nuclear proteins and promotes their subsequent degradation. The heterodimeric ubiquitin-conjugating enzyme UBC13 interacts with RNF8, and they co-localize in the nucleus. RNF8 may regulate mediation of UBC13 polyubiquitylation by elongating the ubiquitin chains. RNF8 also binds to Retinoid X receptor alpha (RXR?, a member of the steroid hormone receptor superfamily. It increases RXR?mediated transactivation of the RXR?responsive element (RXRE) promoter in a dose-dependent manner, suggesting that RNF8 is a regulator of RXR?mediated transcriptional activity.

Gene ID:

9025

Pathways:

Production of Molecular Mediator of Immune Response

#### **Application Details**

**Application Notes:** 

WB 1:300-5000

IHC-P 1:200-400

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months