antibodies -online.com





anti-RLIM antibody (AA 550-600) (Alexa Fluor 350)



Go to Product page

| \sim | | | |
|--------|-----|------|-----|
| | N/P | r\/I | i⊢₩ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | RLIM | |
| Binding Specificity: | AA 550-600 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This RLIM antibody is conjugated to Alexa Fluor 350 | |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |
| Product Details | | |
| Immunogen: | KLH conjugated synthetic peptide derived from human RNF12 | |
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Purification: | Purified by Protein A. | |
| Target Details | | |
| Target: | RLIM | |
| Alternative Name: | RNF12 (RLIM Products) | |
| Background: | Synonyms: RNF12, NY-REN-43, E3 ubiquitin-protein ligase RLIM, LIM domain-interacting RING | |

finger protein, RING finger LIM domain-binding protein, R-LIM, RING finger protein 12, Renal

carcinoma antigen NY-REN-43, RLIM

Background: E3 ubiquitin-protein ligase. Acts as a negative coregulator for LIM homeodomain transcription factors by mediating the ubiquitination and subsequent degradation of LIM cofactors LDB1 and LDB2 and by mediating the recruitment the SIN3a/histone deacetylase corepressor complex. Ubiquitination and degradation of LIM cofactors LDB1 and LDB2 allows DNA-bound LIM homeodomain transcription factors to interact with other protein partners such as RLIM. Plays a role in telomere length-mediated growth suppression by mediating the ubiquitination and degradation of TERF1. By targeting ZFP42 for degradation, acts as an activator of random inactivation of X chromosome in the embryo, a stochastic process in which one X chromosome is inactivated to minimize sex-related dosage differences of X-encoded genes in somatic cells of female placental mammals.

Gene ID:

51132

UniProt:

Q9NVW2

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

| Format: | Liquid | |
|--------------------|--|--|
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |