antibodies -online.com







anti-VGLL4 antibody (AA 220-270) (Biotin)



O -		D	4	page
1-0	$T \cap$	$Pr \cap C$	шет	nane
\cup	LO.	1 100	IUCL	Dauc

()	ve	K\ /		
	\cup	1 V/	-	V۷

Quantity:	100 μL
Target:	VGLL4
Binding Specificity:	AA 220-270
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VGLL4 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human VGLL4	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	VGLL4
Alternative Name:	VGLL4 (VGLL4 Products)
Background:	Synonyms: Transcription cofactor vestigial like protein 4, Vestigial like protein 4, VGL 4, VGLL4, VGLL4_HUMAN.

Background: The transcriptional enhancer factor-1 (TEF-1) family of transcription factors regulate tissue-specific gene expression in muscle and placenta. The mechanism whereby TEF-1 confers tissue specificity depends largely on the interaction of TEF-1 with tissue-specific cofactors. Transcription cofactor Vgl-4 (vestigial-like protein 4) is a 290 amino acid nuclear protein that interacts with TEF-1 and MEF-2. Vgl-4 is the only member of the vestigial-like family that is expressed in heart. Overexpression of Vgl-4 in cardiac myocytes interferes with basal expression and ?-adrenergic receptor-dependent activation of a TEF-1 dependent skeletal ?actin promoter. This suggests that Vgl-4 counteracts ?-adrenergic activation of gene expression in cardiomyocytes. There are two isoforms of Vgl-4 that are produced as a result of alternative splicing events.

Gene ID:

9686

UniProt:

Q14135

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months