

## Datasheet for ABIN1401475

# anti-SERPINA10 antibody (Biotin)



Go to Product page

$\sim$				
( )\	/e	r\/		٨
( ) 1	v C.	ı vı	$\Box$	ΙV

Quantity:	100 μL
Target:	SERPINA10
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SERPINA10 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ZPI/SERPINA10
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

#### **Target Details**

- arget Details	
Target:	SERPINA10
Alternative Name:	SERPINA10 (SERPINA10 Products)
Background:	Synonyms: Protein Z dependent protease inhibitor, Protein Z dependent protease inhibitor
	precursor, Protein Z-dependent protease inhibitor, PZ dependent protease inhibitor, PZ-
	dependent protease inhibitor, PZI, serine or cysteine proteinase inhibitor clade A alpha 1
	antiproteinase antitrypsin member 10 , Serpin A10, Serpin peptidase inhibitor clade A alpha 1

antiproteinase antitrypsin member 10, SERPINA 10, SERPINA10, ZPI, ZPI\_HUMAN.

Background: ZPI, also known as SERPINA10 (serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 10) or PZI, is a 444 amino acid secreted protein that functions as a Protein Z-dependent protease inhibitor. Expressed by the liver, ZPI is secreted into the plasma where, in the presence of calcium, Protein Z and phospholipids, it inhibits the activated pro-coagulation factors X and XI (Factor X and Factor XI). This inhibition helps properly regulate intravenous blood clotting. ZPI, a member of the serpin protein family, contains five potential N-linked glycosylation sites and a tyrosine at position 387 which, when disrupted, renders ZPI inactive. Defects in the gene encoding ZPI may increase susceptibility to venous thrombosis, the formation of blood clots within a vein.

Gene ID:

51156

IHC-P 1:200-400

### **Application Details**

Application Notes: WB 1:300-5000

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months