antibodies .- online.com





anti-RANBP2 antibody (Alexa Fluor 488)



Go to Product page

| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|--------------|--|
| Target: | RANBP2 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RANBP2 antibody is conjugated to Alexa Fluor 488 |
| Application: | Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human RanBP2 | |
|-------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | RANBP2 | |
|-------------------|---|--|
| Alternative Name: | RanBP2 (RANBP2 Products) | |
| Background: | Synonyms: Nuclear pore complex protein Nup358, Nucleoporin 358, Nucleoporin Nup358, NuP358, P270, RAN binding protein 2, Ran-binding protein 2, RANBP2, RBP2_HUMAN, Transformation related protein 2, TRP 1, TRP 2, TRP1, TRP2, 358 kDa nucleoporin, E3 SUMO-protein ligase RanBP2. | |

Background: The nuclear pore complex protein, Ran-binding protein 2 (Ran BP-2 or Nup358), contains four Ran-binding domains. Ran BP-2 is a large scaffold cyclophilin-related protein expressed in photoreceptor cells. Ran BP-2 localization at cytoplasmic fibrils emanating from the nuclear pore complex and interaction with the Ran-GTPase support its role in nucleocytoplasmic transport processes. In humans, the Ran BP-2 gene is partially duplicated in a gene cluster and lies in a hot spot for recombination on chromosome 2q. This genetic heterogeneity renders further significance of this genomic region in human disease due to its possible involvement in genetically linked disorders such as juvenile nephronophthisis, congenital hepatic fibrosis and chorioretinal dysplasia.

Gene ID:

5903

For Research Use only

Pathways:

Carbohydrate Homeostasis, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
| | |

Handling

Restrictions:

| Handling | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |