# antibodies -online.com





# anti-MARCH7 antibody (AA 65-115) (Biotin)



Go to Product page

$\sim$	
( )\/\	rview
$\circ$	1 410 44

Quantity:	100 μL	
Target:	MARCH7	
Binding Specificity:	AA 65-115	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MARCH7 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunoprecipitation (IP)	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MARCH7	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	MARCH7
Alternative Name:	MARCH7 (MARCH7 Products)
Background:	Synonyms: AXO, AXOT, Axotrophin, E3 ubiquitin-protein ligase MARCH7, MARCH VII, MARCH-

VII, March7, MARH7\_HUMAN, Membrane associated RING CH protein VII, Membrane associated ring finger C3HC4 7 E3 ubiquitin protein ligase, membrane associated ring finger C3HC4 7, Membrane associated RING finger protein 7, Membrane-associated RING finger protein 7, Membrane-associated RING-CH protein VII, RING finger protein 177, RNF177. Background: Axotrophin is a stem cell gene that encodes a protein which is involved in T lymphocyte regulation (especially in regulating the proliferation) and leukemia inhibitory factor (LIF) release. LIF is a neuropoietic cytokine that is important for stem cell regulation and thymocyte stimulation. Both Axotrophin and LIF are linked to transplantation intolerance. Axotrophin is also involved in corpus callosum differentiation and may play a role in glial cell line-derived neurotrophic factor (GDNF)-dependent sensory neuron survival in the substantia gelatinosa of the adult spinal cord. Axotrophin is primarily expressed in the hippocampus, cortex, purkinje and granule cells of the cerebellum.

Gene ID: 64844

UniProt: Q9H992

### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

 $IP(1-2 \mu g)$ 

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

1	1 11	١٠
$\vdash$	land	∥n∩
	ıarıa	

Expiry Date:

12 months