antibodies .- online.com





anti-PSMA2 antibody (Biotin)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	PSMA2	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PSMA2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Proteasome 20S alpha 2	
Isotype:	IgG	

Target Details

Cross-Reactivity:

Purification:

Target:	PSMA2	
Abstract:	PSMA2 Products	
Background:	Synonyms: 20S Proteasome 2, HC3, Macropain subunit C3, MU antibody Multicatalytic endopeptidase complex subunit C3, PMSA2, Proteasome prosome macropain subunit alpha	
	type 2, Proteasome alpha 2 subunit, Proteasome component C3, Proteasome subunit alpha	
	type 2, Proteasome subunit alpha type-2, Proteasome subunit HC3, PSC2, PSC3, PSMA 2,	

Human, Mouse, Rat

Purified by Protein A.

Background: The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH . The proteasome has an ATP-dependent proteolytic activity. PSMA2 may have a potential regulatory effect on another component(s) of the proteasome complex through tyrosine phosphorylation.

Gene ID:

5683

Pathways:

Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	