

[Go to Product page](#)

Datasheet for ABIN1402252 **anti-PSMA7 antibody (Alexa Fluor 488)**

Overview

| | |
|--------------|---|
| Quantity: | 100 µL |
| Target: | PSMA7 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PSMA7 antibody is conjugated to Alexa Fluor 488 |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human Proteasome 20S alpha 7 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------|---|
| Target: | PSMA7 |
| Abstract: | PSMA7 Products |
| Background: | Synonyms: C6 antibody HSPC, Proteasome prosome macropain subunit alpha type 7, Proteasome alpha 7 subunit, Proteasome subunit alpha 4, Proteasome subunit alpha type 7, Proteasome subunit alpha type-7, Proteasome subunit RC6 1, Proteasome subunit RC6-1, Proteasome subunit XAPC7, PSA7_HUMAN, PSMA7, RC6 1, XAPC7. |

Target Details

Background: The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH . The proteasome has an ATP-dependent proteolytic activity. Plays an important role in the regulation of cell proliferation or cell cycle control, transcriptional regulation, immune and stress response, cell differentiation, and apoptosis. Interacts with some important proteins involved in transcription factor regulation, cell cycle transition, viral replication and even tumor initiation and progression. Inhibits the transactivation function of HIF-1A under both normoxic and hypoxia-mimicking conditions. The interaction with EMAP2 increases the proteasome-mediated HIF-1A degradation under the hypoxic conditions. Plays a role in hepatitis C virus internal ribosome entry site-mediated translation. Mediates nuclear translocation of the androgen receptor (AR) and thereby enhances androgen-mediated transactivation. Promotes MAVS degradation and thereby negatively regulates MAVS-mediated innate immune response.

Gene ID: 5688

Pathways: [Mitotic G1-G1/S Phases](#), [DNA Replication](#), [Synthesis of DNA](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months