antibodies -online.com





anti-DCAF12 antibody (FITC)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Background:

Quantity:	100 μL	
Target:	DCAF12	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DCAF12 antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human DCAF12/WDR40A	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	DCAF12	
Alternative Name:	DCAF12 (DCAF12 Products)	

 $Order\ at\ www.antibodies-online.com\ |\ www.antiboerper-online.de\ |\ www.anticorps-enligne.fr\ |\ www.antibodies-online.com\ |\ www.antiboerper-online.de\ |\ www.antiboerper-online.d$ International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1402460 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

WDR40A.

Synonyms: cancer/testis antigen 102, Centrosome-related protein TCC52, CT102,

DCA12_HUMAN, DCAF12, DDB1- and CUL4-associated factor 12, TCC52, Testis cancer

centrosome-related protein, WD repeat domain 40A, WD repeat-containing protein 40A,

Background: WD-repeats are motifs that are found in a variety of proteins and are characterized by a conserved core of 40-60 amino acids that commonly form a tertiary propeller structure. While proteins that contain WD-repeats participate in a wide range of cellular functions, they are generally involved in regulatory mechanisms concerning chromatin assembly, cell cycle control, signal transduction, RNA processing, apoptosis and vesicular trafficking. WDR40A (WD-repeat-containing protein 40A), also known as DDB1- CUL4-associated factor 12, is a 453 amino acid cytoplasmic protein that contains four WD-repeats. WDR40A is highly expressed in some cancer cell lines, lung cancer tissues and normal testis. A probable substrate receptor for CUL4-DDB1 E3 ubiquitin-protein ligase complex, WRD40A is suggested to interact with DDB1. WDR40A is encoded by a gene mapping to human chromosome 9p13.3. Human chromosome 9 houses over 900 genes and comprises nearly 4 % of the human genome.

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months