# antibodies -online.com





## anti-ZFPM1 antibody (AbBy Fluor® 555)



$\sim$	+~	$D_{ra} \sim 1$		page
( ¬( )	1()	PIOO	11021	nace

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	ZFPM1	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZFPM1 antibody is conjugated to AbBy Fluor® 555	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FOG1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	ZFPM1	
Alternative Name:	FOG1 (ZFPM1 Products)	
Background:	Synonyms: FOG-1, FOG1_HUMAN, Friend of GATA 1, Friend of GATA protein 1, Friend of GATA1, ZFPM1, Zinc finger protein, Zinc finger protein 89A, Zinc finger protein multitype 1, Zinc finger protein ZFPM1.	
	Background: Transcription regulator that plays an essential role in erythroid and	

#### **Target Details**

megakaryocytic cell differentiation. Essential cofactor that acts via the formation of a heterodimer with transcription factors of the GATA family GATA1, GATA2 and GATA3. Such heterodimer can both activate or repress transcriptional activity, depending on the cell and promoter context. The heterodimer formed with GATA proteins is essential to activate expression of genes such as NFE2, ITGA2B, alpha- and beta-globin, while it represses expression of KLF1. May be involved in regulation of some genes in gonads. May also be involved in cardiac development, in a non-redundant way with ZFPM2/FOG2.

Gene ID:

161882

 $1 \mu g/\mu L$ 

## **Application Details**

Concentration:

Application Notes:	IF(IHC-P) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months