# antibodies -online.com





# anti-SMARCD3 antibody (Alexa Fluor 350)



$\sim$	
( )\/\	rview
$\circ$	1 410 44

Quantity:	100 μL
Target:	SMARCD3
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMARCD3 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SMARCD3
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

# **Target Details**

Target:	SMARCD3
Alternative Name:	SMARCD3 (SMARCD3 Products)
Background:	Synonyms: 60 kDa BRG 1/Brm associated factor subunit C, 60 kDa BRG-1/Brm-associated factor subunit C, BAF60C, BRG1 associated factor 60C, BRG1-associated factor 60C, CRACD3, MGC111010, Rsc6p, Smarcd3, SMRD3_HUMAN, SWI/SNF related matrix associated actin
	dependent regulator of chromatin subfamily d member 3, SWI/SNF-related matrix-associated

actin-dependent regulator of chromatin subfamily D member 3.

Background: SMARCD3, is a member of the SMARCD family and contains one SWIB domain. Two isoforms, isoform 1 and isoform 2 exist due to alternative splicing events. Both isoforms are expressed in placenta, salivary gland, kidney, brain, trachea, uterus, prostate, testis, thyroid, spleen and heart, while isoform 1 is also expressed in adipose tissue and skeletal muscle. Localizing to the nucleus, SMARCD3 is a component of the ATP-dependent chromatin remodeling complex SNF/SWI and is believed to play a role in nucleosome remodeling. SMARCD3 also plays an important role in the regulation of muscle development. In mice, the silencing of the gene en-coding SMARCD3 leads to defects in heart morphogenesis. In addition, both isoforms of SMARCD3 directly interact with and function as coactivators for several transcription factors.

Gene ID:

6604

Pathways:

Regulation of Lipid Metabolism by PPARalpha

### **Application Details**

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months