

Datasheet for ABIN1403065

anti-C9orf152 antibody (Biotin)



\sim	+ ~	רת	$\sim \sim$	uct	00	\sim
(¬()	1()	r	()(1	1101	112	$(1 \leftarrow$

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Overview	
Quantity:	100 μL
Target:	C9orf152
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C9orf152 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human C9orf152
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	C9orf152
Alternative Name:	C9orf152 (C9orf152 Products)
Background:	Synonyms: Chromosome 9 open reading frame 152, Hypothetical protein LOC401546,
	Uncharacterized protein C9orf152, CI152_HUMAN.
	Background: C9orf152 is a 239 amino acid protein encoded by a gene that maps to human
	chromosome 9q31.3. Chromosome 9 consists of about 145 million bases, represents 4 % of

the human genome and encodes nearly 900 genes. Thought to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of BCR-ABL fusion protein often found in leukemias.

Gene ID:

401546

Application Details

Application Notes: WB 1:300-5000

Restrictions:

For Research Use only

IHC-P 1:200-400

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months