antibodies -online.com





anti-ALAS1 antibody (AbBy Fluor® 488)



Go to Product page

()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μL ALAS1 Human, Mouse, Rat	
Target:		
Reactivity:		
Host:	Rabbit	
Clonality:	Polyclonal This ALAS1 antibody is conjugated to AbBy Fluor® 488 Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Conjugate:		
Application:		

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ALAS1/ALAS-H
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	ALAS1
Alternative Name:	ALAS1 (ALAS1 Products)
Background:	Synonyms: mitochondrial, nonspecic, 5 aminolevulinate synthase, 5 aminolevulinate synthase nonspecic mitochondrial, 5 aminolevulinic acid synthase, 5-aminolevulinate synthase, 5-aminolevulinic acid synthase 1, Alas 1, ALAS 3, ALAS H, ALAS N, ALAS-H, alaS1, ALAS3, ALASH, Aminolevulinate delta synthase 1, Aminolevulinic acid synthase 1, Delta ALA

synthetase, Delta aminolevulinate synthase, Delta-ALA synthase 1, Delta-aminolevulinate
synthase 1, HEM1_HUMAN, MIG 4, MIG4, Migration inducing protein 4.

Background: 5-aminolevulinate synthase 1 (ALAS-H) and 2 (ALAS-E) are two isoforms of ALAS, an enzyme catalyzing the first step of the heme biosynthetic pathway in mammals. The erythroid-specific isoenzyme, ALAS-E, regulates the first step of hematopoietic cell differentation and iron metabolism in the liver. ALAS-H is a housekeeping protein which mediates synthesis of early heme in the mitochondria of most cells. Succinyl CoA associates with ALAS-E in protein conformation change and translocation of ALAS-E into the mitochondria and does not interact with ALAS-H. The ALAS-E 5'-flanking region contains binding sites for nuclear activators such as GATA-1, NF-E2 and EKLF. Since the ALAS gene maps to the X chromosome, mutation of the gene leads to the pyridoxine-refractory X-linked sideroblastic anemia.

Gene ID: 211

Pathways: Regulation of Lipid Metabolism by PPARalpha

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months