# antibodies -online.com





## anti-GLT8D1 antibody (Biotin)



Go to Product page

| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:    | 100 μL   |  |
|--------------|--|--|
| Target:      | GLT8D1   |  |
| Reactivity:  | Human, Mouse, Rat  |  |
| Host:        | Rabbit   |  |
| Clonality:   | Polyclonal   |  |
| Conjugate:   | This GLT8D1 antibody is conjugated to Biotin                                       |  |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |  |

### **Product Details**

| Immunogen:        | KLH conjugated synthetic peptide derived from human GLT8D1 |  |
|-------------------|--|--|
| Isotype:          | IgG  |  |
| Cross-Reactivity: | Human, Mouse, Rat  |  |
| Purification:     | Purified by Protein A.                                     |  |

## **Target Details**

| Target:           | GLT8D1   |  |
|-------------------|--|--|
| Alternative Name: | GLT8D1 (GLT8D1 Products)   |  |
| Background:       | Synonyms: MGC94018, Da2 24, GALA4A, Glycosyltransferase 8 domain containing 1, Glycosyltransferase 8 domain-containing protein 1, Glycosyltransferase AD 017, MSTP139, |  |
|                   | GL8D1_HUMAN.   |  |
|                   | Background: GLT8D1 is a 371 amino acid single-pass type II transmembrane protein that is   |  |

#### **Target Details**

expressed by a gene residing on human chromosome 3. Chromosome 3 is made up of about 214 million bases encoding over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci. Key tumor suppressing genes on chromosome 3 include those that encode the apoptosis mediator RASSF1, the cell migration regulator HYAL1 and the angiogenesis suppressor SEMA3B. Marfan Syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth Disease are a few of the numerous genetic diseases associated with chromosome 3. There are two isoforms of GLT8D1 that are produced as a result of alternative splicing events.

Gene ID:

55830

For Research Use only

## **Application Details**

| Application Notes: | WB 1:300-5000   |
|--------------------|-----------------|
|                    | IHC-P 1:200-400 |

Restrictions:

Handling

| - Idridiirig       |  |
|--------------------|--|
| Format:            | Liquid   |
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C for 12 months.  |
| Expiry Date:       | 12 months  |