

[Go to Product page](#)

## Datasheet for ABIN1403647 **anti-PIBF1 antibody (Biotin)**

### Overview

Quantity:	100 µL
Target:	PIBF1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIBF1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PIBF1/C13orf24
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### Target Details

Target:	PIBF1
Alternative Name:	PIBF1 ( <a href="#">PIBF1 Products</a> )
Background:	Synonyms: PIBF 1, PIBF, PIBF1, PIBF-1, Progesterone immunomodulatory binding factor 1, Progesterone induced blocking factor 1, Progesterone-Induced Blocking Factor 1, RP11 505F3.1, PIBF1_HUMAN, C13orf24. Background: PIBF is synthesized during pregnancy in response to progesterone by

## Target Details

progesterone receptor-positive T lymphocytes (mostly gamma-delta T cells). In the presence of PIBF, natural killer (NK) cells inhibit the release of perforin from storage granules and therefore fail to lyse target cells. In humans, the amount of cells that express PIBF is significantly higher in healthy pregnant women than in women at risk for premature pregnancy termination. Full-length PIBF is associated with the nucleus, whereas secretion of shorter forms is induced by activation of the cell. Research suggests that PIBF functions as a transcription factor in its full-length form, while smaller forms may act as cytokines. The PIBF gene encodes a deduced hydrophilic 757-amino acid alpha-helical protein with an N-terminal signal sequence, a leucine zipper motif, a basic zipper sequence, a PEST sequence, a nuclear localization signal, an endoplasmic reticulum membrane retention signal, and many presumed N-glycosylation and phosphorylation sites.

Gene ID: 10464

## Application Details

Application Notes: WB 1:300-5000  
IHC-P 1:200-400

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months