# antibodies -online.com





# anti-Retinoic Acid Induced 12 (RAI12) antibody (AbBy Fluor® 647)



Go to Product page

0	1 /	0	K	/ 1	$\sim$	1 A /
	1//	$\vdash$	I \	71	$\vdash$	<b>\/\/</b>

Quantity:	100 μL
Target:	Retinoic Acid Induced 12 (RAI12)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	AbBy Fluor® 647
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DERP6/C17orf81	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

# Target Details

Target:	Retinoic Acid Induced 12 (RAI12)	
Alternative Name:	DERP6 (RAI12 Products)	
Background:	Synonyms: MST071, MSTP071, chromosome 17 open reading frame 81, Dermal papilla derived	
	protein 6, dermal papilla-derived protein 6, DERP6, HSPC002, S phase 2 protein, ELP5_HUMAN.	
	Background: DERP6, is a 316 amino acid protein that localizes to the cytoplasm and exists as	
	multiple alternatively spliced isoforms. Expressed ubiquitously with highest expression in liver,	

heart, testis, brain and skeletal muscle, DERP6 is thought to be involved in p53-mediated transcriptional regulation. The gene encoding DERP6 maps to human chromosome 17, which comprises over 2.5 % of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes.

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	