antibodies -online.com







anti-C17orf82 antibody (Biotin)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	C17orf82	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C17orf82 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C17orf82	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

Target Details

Target:	C17orf82	
Alternative Name:	C17orf82 (C17orf82 Products)	
Background:	Synonyms: C17orf82, Chromosome 17 open reading frame 82, CQ082_HUMAN, Putative uncharacterized protein C17orf82. Background: C17orf82 is a 251 amino acid protein that is encoded by a gene mapping to	
	human chromosome 17. Chromosome 17 makes up over 2.5 $\%$ of the human genome with	

about 81 million bases encoding over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes. Chromosome 17 is also linked to neurofibromatosis, a condition characterized by neural and epidermal lesions, and dysregulated Schwann cell growth. Alexander disease, Birt-Hogg-Dube syndrome and Canavan disease are also associated with chromosome 17.

Gene ID:

388407

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months